

Canonical forms of $2 \times 3 \times 3$ tensors over the real field, algebraically closed fields, and finite fields.

Michel Lavrauw* and John Sheekey†

February 11, 2015

Abstract

We classify the orbits of elements of the tensor product spaces $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$ for all finite; real; and algebraically closed fields under the action of two natural groups. The result can also be interpreted as the classification of the orbits in the 17-dimensional projective space of the Segre variety product of a projective line and two projective planes. This extends the classification of the orbits in the 7-dimensional projective space of the Segre variety product of three projective lines [9]. The proof is geometric in nature, relies on properties of the Segre embedding, and uses the terminology of projective spaces.

1 Introduction

1.1 Motivation and main results

In this article we study orbits of elements of tensor product spaces $V_1 \otimes V_2 \otimes V_3$ under the action of two natural groups. This problem can be studied either

*The research of the first author was supported by the Fund for Scientific Research - Flanders (FWO) and by a Progetto di Ateneo from Università di Padova (CPDA113797/11).

†The second author acknowledges the support of the Fund for Scientific Research - Flanders (FWO).

in the vector space $V = V_1 \otimes V_2 \otimes V_3$ or the projective space $\text{PG}(V_1 \otimes V_2 \otimes V_3)$. In this article we will present results using vector space terminology. However we will utilise some terminology and concepts from projective geometry in the proofs.

We consider orbits under the action of the groups G and H , where G is the stabiliser in $\text{GL}(V)$ of the set of *fundamental tensors* (see Section 2 for definitions), and H is the subgroup $\text{GL}(V_1) \times \text{GL}(V_2) \times \text{GL}(V_3)$ of G .

These orbits are interesting not only for their own sake, being as they are fundamental objects in both algebra and geometry. They also have many important applications: see for example [6, Chapter 1]. Furthermore, tensors over finite fields have applications in the study of *finite semifields* (that is, nonassociative division algebras), an important and active research topic in finite geometry, see e.g. [8]. A deeper understanding of the geometry of three-fold tensor product spaces is crucial to a better structural insight into the set of the isotopism classes of finite semifields, see [7]. This paper provides a further step in this direction by classifying the orbits in $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^3$ and $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$, in the case that $\mathbb{F}$ is a finite field, the real field, or algebraically closed. The proof uses ideas from [7], [9], and [10], and gives geometric characterisations of each orbit.

Main Theorem. *If $\mathbb{F}$ is a finite field, then there are precisely 21 H -orbits and 18 G -orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$. For any algebraically closed field $\mathbb{F}$, there are precisely 18 H -orbits and 15 G -orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$. There are precisely 20 H -orbits and 17 G -orbits of tensors in $\mathbb{R}^2 \otimes \mathbb{R}^3 \otimes \mathbb{R}^3$.*

This result can be used to fully classify all orbits of subspaces of $\text{PG}(\mathbb{F}^2 \otimes \mathbb{F}^3)$, and hence all orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^r$ for all positive integers r . This will be treated by the authors in the forthcoming paper [11].

1.2 Historical context

The orbits under the actions of G and H have been studied mostly over algebraically closed fields, see for example [6]. The classification of G - (resp. H -)orbits of $3 \times 3 \times 3$ tensors over the complex numbers has been done in [14] (resp. [12]), and the result for complex $2 \times 3 \times 3$ tensors can be obtained from these. However the techniques employed therein (classification of complex ternary cubics in [14], topology in [12]) are not applicable over finite fields.

The classification of $2 \times m \times n$ tensors can be approached using the Weierstrass-Kronecker theory of pencils (that is, $m \times n$ matrices with polynomial entries linear in an indeterminate λ), see for example [5]. From each tensor we can obtain a pencil in a canonical way. However, the classification of pencils does not imply a classification of G - or H -orbits, as equivalent tensors can give rise to inequivalent pencils. Furthermore, this approach does not give any geometric insight to the nature of these orbits, and does not extend well to larger order tensors.

This approach was used in [15] to give canonical forms for $2 \times 3 \times 3$ tensors over fields of odd prime order $p > 3$. However the forms are presented without proof. For example, there is no proof in the literature that the canonical forms $M_2(x)$ and $M_8(x)$ in [15, Section 7] are indeed the unique orbits with the relevant invariant factors (and in fact these forms are not valid over fields of even characteristic).

Computational and theoretical results over small fields and small dimensions have been obtained, for example [1], [2], [13]. A geometric classification for the case $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^2$ was given by the authors in [9].

The main result of this paper is to classify orbits of $2 \times 3 \times 3$ tensors over all finite fields, and geometrically characterise each orbit. As our approach will be elementary and mostly field-independent, we will also re-obtain the classification over the complex numbers and prime fields, and moreover over any algebraically closed field, and the real numbers.

2 Definitions and preliminary results

Let $V = \bigotimes_{i=1}^r V_i$, where $V_1, \dots, V_r$ are finite dimensional vector spaces over some field $\mathbb{F}$, with $\dim V_i = n_i < \infty$. The set of *fundamental tensors* is the set $\{v_1 \otimes v_2 \otimes \dots \otimes v_r : v_i \in V_i \setminus \{0\}\}$. Projectively, this corresponds to the *Segre variety* $S_{n_1, n_2, \dots, n_r}(\mathbb{F})$, that is the image of a *Segre embedding* $\sigma_{n_1, \dots, n_r}$ defined as

$$\begin{aligned} \sigma_{n_1, \dots, n_r} : \text{PG}(V_1) \times \text{PG}(V_2) \times \dots \times \text{PG}(V_r) &\rightarrow \text{PG}(V) \\ : (\langle v_1 \rangle, \langle v_2 \rangle, \dots, \langle v_r \rangle) &\mapsto \langle v_1 \otimes v_2 \otimes \dots \otimes v_r \rangle. \end{aligned}$$

The *rank* of a tensor A , denoted $\text{rk}(A)$, is defined to be the minimum number k such that there exist fundamental tensors $\alpha_1, \dots, \alpha_k$ with $A \in \langle \alpha_1, \dots, \alpha_k \rangle$.

The fundamental tensors hence correspond precisely to the set of *rank one tensors*. When $r = 2$, that is $V = V_1 \otimes V_2$, the tensor rank corresponds to the usual matrix rank. We define the rank of a projective point in $\text{PG}(V)$ to be the rank of any corresponding tensor.

The setwise stabilizer in $\text{GL}(V)$ of the set of rank one tensors in $V = V_1 \otimes V_2 \otimes V_3$ will be denoted by G . Clearly the rank of a tensor is G -invariant. The group $\text{GL}(V_1) \times \text{GL}(V_2) \times \text{GL}(V_3)$ defines a subgroup of G , which we will denote by H , via the action defined by

$$(g_1, g_2, g_3) : v_1 \otimes v_2 \otimes v_3 \mapsto v_1^{g_1} \otimes v_2^{g_2} \otimes v_3^{g_3}.$$

Note that H is not necessarily equal to G , for example if $V_1 \neq V_2 = V_3$, then $G \cong \text{GL}(V_1) \times (\text{GL}(V_2) \wr \text{Sym}(2))$. The group $\text{Sym}(2)$ arises from the map T defined by $T : v_1 \otimes v_2 \otimes v_3 \mapsto v_1 \otimes v_3 \otimes v_2$.

For $A \in V_1 \otimes V_2 \otimes V_3$ we define the *first contraction space* of A as the following subspace of $V_2 \otimes V_3$:

$$A_1 := \langle w_1^\vee(A) : w_1^\vee \in V_1^\vee \rangle,$$

where $V_1^\vee$ denotes the dual space of V_1 , and the *contraction* $w_1^\vee(A)$ is defined by its action on the fundamental tensors

$$w_1^\vee(v_1 \otimes v_2 \otimes v_3) = w_1^\vee(v_1)v_2 \otimes v_3. \quad (1)$$

Similarly we define the *second* and *third contraction space*, and denote these by A_2 and A_3 respectively. We will consider the projective subspaces $\text{PG}(A_i)$ of $\text{PG}(V_j \otimes V_k)$, where $j < k$ and $\{i, j, k\} = \{1, 2, 3\}$.

The setwise stabilizer of the set of rank one tensors in the contracted space $V_j \otimes V_k$ will be denoted by G_i , where $j < k$ and $\{i, j, k\} = \{1, 2, 3\}$, and the subgroups $\text{GL}(V_j) \times \text{GL}(V_k)$ of G_i by H_i . For example, with $V_1 = \mathbb{F}^2$, $V_2 = V_3 = \mathbb{F}^n$, $n > 2$, the group $G_1 \cong \text{GL}(n, q) \wr \text{Sym}(2) > H_1$, while $G_2 = H_2$ and $G_3 = H_3$.

The following elementary lemmas will prove useful in the classification to follow.

Lemma 2.1. *If $A, B \in V_1 \otimes V_2 \otimes V_3$ then the following are equivalent:*

- (i) $A \sim_H B$;

(ii) $\forall i \ A_i \sim_{H_i} B_i$;

(iii) $\exists i \ A_i \sim_{H_i} B_i$.

Proof: Suppose $A_1 \sim_{H_1} B_1$, i.e. suppose there exists $(h_2, h_3) \in H_1$ such that $A_1^{(h_2, h_3)} = B_1$. Let $A = \sum a_{ijk} u_i \otimes v_j \otimes w_k$, where the u_i 's, v_i 's, w_i 's form a basis of V_1, V_2, V_3 , respectively. Then

$$A_1 = \langle \sum a_{ijk} v_j \otimes w_k : i = 1, \dots, \dim V_1 \rangle, \quad (2)$$

and

$$B_1 = A_1^{(h_2, h_3)} = \langle \sum a_{ijk} v_j^{h_2} \otimes w_k^{h_3} : i = 1, \dots, \dim V_1 \rangle. \quad (3)$$

This implies that $B = \sum b_i u'_i \otimes (a_{ijk} v_j^{h_2} \otimes w_k^{h_3})$ where $b_i \in \mathbb{F}$, and the u'_i 's form a basis. Hence $B = \sum u_i^{h_1} \otimes (a_{ijk} v_j^{h_2} \otimes w_k^{h_3}) = A^{(h_1, h_2, h_3)}$, where $\text{GL}(V_1) \ni h_1$ maps u_i to $b_i u'_i$ for $b_i \neq 0$ and u_i to u'_i otherwise. This proves $(iii) \Rightarrow (i)$. The proof of $(i) \Rightarrow (ii)$ is similar, and $(ii) \Rightarrow (iii)$ is trivial. $\square$

Corollary 2.2. *Suppose $V_1 \neq V_2 = V_3$. Then two tensors A and B are G -equivalent if and only if A_1 is H_1 -equivalent to one of $\{B_1, B_1^T\}$, if and only if A_1 is G_1 -equivalent to B_1 .*

Suppose $V_2 = V_3 = \mathbb{F}^n$, and let $\{e_1, \dots, e_n\}$ be the standard basis for $\mathbb{F}^n$. Then $V_2 \otimes V_3 \simeq M_n(\mathbb{F})$, and we can choose the isomorphism such that the tensor $x := \sum_{i=1}^n e_i \otimes e_i$ corresponds to the identity matrix. Then the stabiliser of x in H_1 is given by $N := \{(g, g^{-1}) : g \in \text{GL}(n, \mathbb{F})\}$.

Lemma 2.3. *If S_U denotes the stabiliser in $\text{GL}(V)$ of a k -dimensional subspace U of a vector space V . Then $\{g^{-1} : g \in S_U\}$ is the stabiliser in $\text{GL}(V)$ of a subspace of codimension k .*

Proof: Let V_1 denote the n -dimensional vector space of row vectors and V_2 the n -dimensional vector space of column vectors over the field $\mathbb{F}$. Then $\text{GL}(V_1) = \text{GL}(V_2)$ with the action from the right on V_1 and from the left on V_2 . Let U be a subspace of V_1 and define $U^\perp := \{y \in V_2 : \forall u \in U : uy = 0\}$. Suppose $A \in \text{GL}(V_1)$ is in the stabiliser S_U in $\text{GL}(V_1)$ of U . Then $\forall u \in U, uA^{-1} \in U$ and hence $uA^{-1}y = 0, \forall y \in U^\perp$. But this means that $A^{-1}U^\perp \subseteq U^\perp$. It follows that $\{A^{-1} : A \in S_U\}$ is the stabiliser of $U^\perp$ in $\text{GL}(V_2)$. $\square$

Lemma 2.4. *A rank two point x in $\langle S_{n_1, n_2}(\mathbb{F}) \rangle$ is contained in a unique solid which intersects $S_{n_1, n_2}(\mathbb{F})$ in a subvariety $Q(x)$ equivalent to a Segre variety $S_{2,2}(\mathbb{F})$.*

Proof: Consider a point x of rank two, i.e. a point $x \in \langle y, z \rangle$ where $y, z \in S_{n_1, n_2}(\mathbb{F})$, $y \neq x \neq z$. If $y = \sigma_{n_1, n_2}(y_1 \times y_2)$ and $z = \sigma_{n_1, n_2}(z_1 \times z_2)$, then $Q_{y,z} := \sigma_{n_1, n_2}(\langle y_1, z_1 \rangle \times \langle y_2, z_2 \rangle)$ is a Segre variety equivalent to $S_{2,2}(\mathbb{F})$ contained in $S_{n_1, n_2}(\mathbb{F})$, and $x \in \langle Q_{y,z} \rangle$. Any subvariety of $S_{n_1, n_2}(\mathbb{F})$ equivalent to $S_{2,2}(\mathbb{F})$ is of the form $Q_{y,z}$ for some $y, z \in S_{n_1, n_2}(\mathbb{F})$, and two solids spanned by such subvarieties $Q_{y,z} = \sigma_{n_1, n_2}(\langle y_1, z_1 \rangle \times \langle y_2, z_2 \rangle)$ and $Q_{y',z'} = \sigma_{n_1, n_2}(\langle y'_1, z'_1 \rangle \times \langle y'_2, z'_2 \rangle)$ intersect in the subspace spanned by the subvariety $\sigma_{n_1, n_2}(w_1 \times w_2)$, where $w_1 = \langle y_1, z_1 \rangle \cap \langle y'_1, z'_1 \rangle$ and $w_2 = \langle y_2, z_2 \rangle \cap \langle y'_2, z'_2 \rangle$. It follows that the intersection of two different solids $\langle Q_{y,z} \rangle$ and $\langle Q_{y',z'} \rangle$ is either a point on $S_{n_1, n_2}(\mathbb{F})$ or a line on $S_{n_1, n_2}(\mathbb{F})$, and hence cannot contain a point of rank two. This implies that $Q(x) := Q_{y,z}$ is uniquely defined by x . $\square$

Definition 2.5. *The (i -th) rank distribution $r_i(A)$ of a tensor A is defined to be the tuple whose j -th entry is the number of rank j points in the i -th contraction space $\text{PG}(A_i)$.*

Note that if A and B are H -equivalent, then $r_i(A) = r_i(B)$ for each i . The converse does not necessarily hold. Also note that the corresponding statement for G -equivalence is false.

3 Orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$

In this section we study the orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$ in terms of their first contraction spaces, by considering the corresponding projective subspaces of $\text{PG}(\mathbb{F}^3 \otimes \mathbb{F}^3) \simeq \text{PG}(8, \mathbb{F})$. By Lemma 2.1 and its corollary, the H_1 -orbits (resp. G_1 -orbits) of these subspaces are in one-to-one correspondence with the H -orbits (resp. G -orbits) of nonzero tensors. The proof itself contains a geometric characterisation of each of the orbits. A nonzero tensor in $\mathbb{F}^3 \otimes \mathbb{F}^3$ (and similarly a point in $\text{PG}(\mathbb{F}^3 \otimes \mathbb{F}^3)$) is called *singular* if it has rank at most two.

3.1 The trivial cases $\mathbf{o}_0, \mathbf{o}_1, \mathbf{o}_2, \mathbf{o}_3$

The zero vector forms an H -orbit, which we denote by $\mathbf{o}_0$. The next trivial cases occur when $\text{PG}(A_1)$ is a point, and this gives rise to three further orbits: $\mathbf{o}_1, \mathbf{o}_2, \mathbf{o}_3$. Namely, if $A \in \mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$, $A \neq 0$, with first contraction space A_1 of dimension one, then A is of the form $x \otimes \tau$, with $x \in \mathbb{F}^2$ and $\tau \in \mathbb{F}^3 \otimes \mathbb{F}^3$. Two such nonzero tensors $A = x \otimes \tau$ and $A' = x' \otimes \tau'$ are G -equivalent if and only if τ and τ' are G_1 -equivalent. Therefore, it is clear that there are three orbits on the nonzero vectors in $\{A \in \mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3 : \dim(A_1) = 1\}$. Call these orbits $\mathbf{o}_1, \mathbf{o}_2$ and $\mathbf{o}_3$, where the subscript i corresponds to the rank of the 2-fold tensor, i.e. $A \in \mathbf{o}_i$ if and only if $\text{rk}(A_1) = i$.

3.2 Lines containing at least one singular point

Next we consider tensors $A \in \mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$ whose first contraction space A_1 is two-dimensional and contains a singular tensor. This means that A_1 determines a line $\text{PG}(A_1)$ containing a singular point.

Theorem 3.1. *Let Ω denote the set of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$ whose first contraction space is two-dimensional and contains at least one singular tensor. If $\mathbb{F}$ is finite, or $\mathbb{F}$ is real closed, there are 16 H -orbits and 13 G -orbits on Ω . If $\mathbb{F}$ is algebraically closed, there are 14 H -orbits and 11 G -orbits on Ω .*

Proof: We will consider the orbits of points and lines in $\text{PG}(\mathbb{F}^3 \otimes \mathbb{F}^3)$ under the action induced by H_1 . We divide the proof into subcases by considering the rank distribution. For a tensor $A \in \mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$ we denote the rank distribution $r_1(A)$ of A by $[a, b, c]$.

Let p_1, p_2, p_3, p_4 be the standard frame of $\text{PG}(\mathbb{F}^3)$ corresponding to the basis e_1, e_2, e_3 of $\mathbb{F}^3$, and recall the Segre embedding

$$\sigma_{3,3} : \text{PG}(\mathbb{F}^3) \times \text{PG}(\mathbb{F}^3) \rightarrow S_{3,3}(\mathbb{F}) \subset \text{PG}(\mathbb{F}^3 \otimes \mathbb{F}^3). \quad (4)$$

Case (1): $\text{PG}(A_1)$ is a line, with $a \geq 2$; $\mathbf{o}_4, \mathbf{o}_4^T, \mathbf{o}_5$.

We are supposing $a \geq 2$, i.e. $\text{PG}(A_1)$ has at least two points of rank 1. The lines contained in $S_{3,3}(\mathbb{F})$ are either of the form $\sigma_{3,3}(p_1 \times \langle p_1, p_2 \rangle)$ or

$\sigma_{3,3}(\langle p_1, p_2 \rangle \times p_1)$ and they form two orbits under the action induced by H_1 . Clearly these are G_1 -equivalent under the action induced by the element $T : v_2 \otimes v_3 \mapsto v_3 \otimes v_2$, and so we denote them by $\mathbf{o}_4$ and $\mathbf{o}_4^T$ respectively. In the case $\mathbb{F} = \mathbb{F}_q$, these lines have rank distribution $[q + 1, 0, 0]$.

The lines not contained in $S_{3,3}(\mathbb{F})$ have $a \leq 2$, since $S_{3,3}(\mathbb{F})$ is the intersection of quadrics. Suppose $a = 2$. Then $\text{PG}(A_1)$ is a two-secant to $S_{3,3}(\mathbb{F})$ and hence determines a unique subvariety $S_{2,2}(\mathbb{F})$. Each such subvariety is equivalent to $\sigma_{3,3}(\langle p_1, p_2 \rangle \times \langle p_1, p_2 \rangle)$, and its stabiliser inside the group induced by H_1 acts 3-transitively on the points of $\langle p_1, p_2 \rangle$ in each factor. Therefore, the set of two-secants forms one orbit under the group induced by H_1 . All other points of $\text{PG}(A_1)$ have rank two and hence if $\mathbb{F} = \mathbb{F}_q$, the line $\text{PG}(A_1)$ has rank distribution $[2, q - 1, 0]$. We denote this orbit by $\mathbf{o}_5$.

Case (2): $\text{PG}(A_1)$ is a line, with $a = 1$; $\mathbf{o}_6, \mathbf{o}_7, \mathbf{o}_7^T, \mathbf{o}_8, \mathbf{o}_9$.

Next suppose that $a = 1$ and $b \geq 1$, i.e. $\text{PG}(A_1)$ is tangent to $S_{3,3}(\mathbb{F})$ and contains at least one point of rank 2. Assume $\text{PG}(A_1) = \langle x_1, x_2 \rangle$, with x_1 of rank one and x_2 of rank two. We consider three cases depending on the position of x_1 with respect to $Q(x_2)$.

$\mathbf{o}_6$. If x_1 is contained in $Q(x_2)$ then $\text{PG}(A_1)$ is a tangent line to $Q(x_2)$ in $\langle Q(x_2) \rangle$ and it follows from [9] that these lines form one orbit under the group induced by H_1 . This orbit has rank distribution $[1, q, 0]$ in case $\mathbb{F} = \mathbb{F}_q$.

$\mathbf{o}_7$. If x_1 is not contained in $Q(x_2)$, but is contained in a subvariety $S_{2,3}(\mathbb{F})$ of $S_{3,3}(\mathbb{F})$ containing $Q(x_2)$, then w.l.o.g. we may assume that $\text{PG}(A_1) = \langle x_1, x_2 \rangle$ is contained in $\text{PG}(\langle e_1, e_2 \rangle \otimes \mathbb{F}^3)$, where $x_2 = \langle e_1 \otimes e_1 + e_2 \otimes e_2 \rangle$ and $x_1 = \langle e_1 \otimes e_3 \rangle$, since the subgroup of the group induced by H_1 stabilising $\sigma_{3,3}(\langle p_1, p_2 \rangle \times \text{PG}(\mathbb{F}^3))$ acts transitively on frames of both factors. Note that the dimension of the third contraction space A_3 is three, since otherwise we would be in the previous case $\mathbf{o}_6$. In the finite case $\mathbb{F} = \mathbb{F}_q$, the line $\text{PG}(A_1)$ has rank distribution $[1, q, 0]$.

$\mathbf{o}_7^T$. If x_1 is not contained in $Q(x_2)$, but is contained in a subvariety $S_{3,2}(\mathbb{F})$ of $S_{3,3}(\mathbb{F})$ containing $Q(x_2)$, repeating the argument for $\mathbf{o}_7$ we see that we get

a single orbit, which is G_1 -equivalent to o_7 , and also has rank distribution $[1, q, 0]$. Note that o_7 and o_7^T cannot be H_1 -equivalent, for then $\text{PG}(A_1)$ would be contained in some $\langle S_{2,2}(\mathbb{F}) \rangle$.

o₈. If x_1 is not contained in a subvariety $S_{2,3}(\mathbb{F})$ or $S_{3,2}(\mathbb{F})$ of $S_{3,3}(\mathbb{F})$ containing $Q(x_2)$, then, again using the transitivity of the group induced by H_1 on frames of both factors, w.l.o.g. we have $x_1 = \langle e_1 \otimes e_1 \rangle$ and $x_2 = \langle e_2 \otimes e_2 + e_3 \otimes e_3 \rangle$. If $\mathbb{F} = \mathbb{F}_q$ then $\text{PG}(A_1)$ has rank distribution $[1, 1, q - 1]$.

o₉. This leaves only one more possibility for tangent lines, namely when $b = 0$ and $a = 1$. An example of such a line is $\langle x_1, x_2 \rangle$ with $x_1 = \langle e_1 \otimes e_3 \rangle$ and $x_2 = \langle e_1 \otimes e_1 + e_2 \otimes e_2 + e_3 \otimes e_3 \rangle$. We show that also these lines form one orbit. For suppose $\text{PG}(A_1)$ is a line containing a point x_1 of rank one and a point x_2 of rank 3. Then w.l.o.g. we may assume that $x_2 = \langle e_1 \otimes e_1 + e_2 \otimes e_2 + e_3 \otimes e_3 \rangle$ and $x_1 = \langle x' \otimes x'' \rangle$ with $x' = ue_1 + ve_2 + we_3$ for some $u, v, w \in \mathbb{F}$ with $u \neq 0$. Now x_2 can also be written as

$$x_2 = \langle (ue_1 + ve_2 + we_3) \otimes \frac{e_1}{u} + e_2 \otimes (e_2 - \frac{v}{u}e_1) + e_3 \otimes (e_3 - \frac{w}{u}e_1) \rangle.$$

After an appropriate base change we see that up to H_1 -equivalence we may assume $x_1 = e_1 \otimes (u'e_1 + v'e_2 + w'e_3)$, for some $u', v', w' \in \mathbb{F}$ with $w' \neq 0$, and $x_2 = e_1 \otimes e_1 + e_2 \otimes e_2 + e_3 \otimes e_3$. Now note that necessarily $u' = 0$, since otherwise the line $\langle x_1, x_2 \rangle$ would contain the rank two point $\langle x_2 - u'x_1 \rangle$ in contradiction with our hypothesis $b = 0$. So w.l.o.g. we have $x_1 = e_1 \otimes (v'e_2 + w'e_3)$. Finally, since x_2 can be rewritten as

$$e_1 \otimes e_1 + (e_2 - \frac{v'}{w'}e_3) \otimes e_2 + \frac{e_3}{w'} \otimes (v'e_2 + w'e_3),$$

we see that the line $\langle x_1, x_2 \rangle$ is equivalent to $\langle e_1 \otimes e_3, e_1 \otimes e_1 + e_2 \otimes e_2 + e_3 \otimes e_3 \rangle$, as claimed. We denote this orbit by **o₉**. If $\mathbb{F} = \mathbb{F}_q$ these lines have rank distribution $[1, 0, q]$.

The orbits $o_1, \dots, o_9$ are the only orbits on lines with rank distribution $[a, b, c]$ with $a \geq 1$. So from now on we assume $a = 0$.

Case (3): $\text{PG}(A_1)$ is a line, with $a = 0$, $b \geq 2$; **o₁₀**, **o₁₁**, **o₁₁^T**, **o₁₂**, **o₁₃**, **o₁₄**.

$\mathbf{o}_{10}$. Our next orbit is inherited from the $2 \times 2 \times 2$ case; it consists of lines which are contained in a three-dimensional subspace spanned by a subvariety $S_{2,2}(\mathbb{F})$ of $S_{3,3}(\mathbb{F})$ but disjoint from this subvariety. These corresponds to the orbit of nonsingular points in the Segre variety product of three projective lines, see [9]. When $\mathbb{F} = \mathbb{F}_q$ or $\mathbb{F} = \mathbb{R}$, by [9] these lie in one orbit which we denote by $\mathbf{o}_{10}$. The rank distribution for $\mathbb{F} = \mathbb{F}_q$ is $[0, q+1, 0]$, i.e. *constant rank two*. Note that this orbit is empty if $\mathbb{F}$ is algebraically closed.

$\mathbf{o}_{11}$. Another orbit containing constant rank two lines, consists of the lines disjoint from $S_{3,3}(\mathbb{F})$ which are contained in a five-dimensional space spanned by some subvariety $S_{2,3}(\mathbb{F})$, but not any $S_{2,2}(\mathbb{F})$. We claim that all these lines are equivalent to the line $\langle e_1 \otimes e_1 + e_2 \otimes e_2, e_1 \otimes e_2 + e_2 \otimes e_3 \rangle$. Suppose $\langle x_1, x_2 \rangle$ is a constant rank two line contained in a subvariety $S_{2,3}(\mathbb{F})$. Then w.l.o.g. we may assume that $\langle x_1, x_2 \rangle$ is contained in $\text{PG}(\langle e_1, e_2 \rangle \otimes \mathbb{F}^3)$. It follows that $Q(x_1)$ and $Q(x_2)$ meet in a line ℓ , and again w.l.o.g. we may assume $\ell = \sigma_{3,3}(\langle e_1, e_2 \rangle \otimes \langle e_2 \rangle)$. Each plane $\pi_i = \langle x_i, \ell \rangle$, $i = 1, 2$, intersects the quadric $Q(x_i)$ in two lines: the line ℓ and another line, say ℓ_i . If ℓ_1 and ℓ_2 have a point in common, then the plane $\langle \ell_1, \ell_2 \rangle$ is contained in the Segre variety $S_{3,3}(\mathbb{F})$. In this case the three-dimensional space $\langle \ell, \ell_1, \ell_2 \rangle$ contains the plane $\langle \ell_1, \ell_2 \rangle$ as well as the line $\langle x_1, x_2 \rangle$. But then $\langle x_1, x_2 \rangle$ meets $\langle \ell_1, \ell_2 \rangle$ contradicting our hypothesis that $\langle x_1, x_2 \rangle$ is a constant rank two line. Therefore $\ell_1 \cap \ell_2 = \emptyset$. W.l.o.g. we may assume that $y_1 = \ell \cap \ell_1 = \langle e_1 \otimes e_2 \rangle$ and $y_2 = \ell \cap \ell_2 = \langle e_2 \otimes e_2 \rangle$. Now considering the intersection point $\langle e_1 \otimes \alpha \rangle$ of the line ℓ_1 with the line $\langle x_1, y_2 \rangle$ and the intersection point $\langle e_2 \otimes \beta \rangle$ of the line ℓ_2 with the line $\langle x_2, y_1 \rangle$ we have that $x_1 = \langle ue_1 \otimes \alpha + e_2 \otimes e_2 \rangle$ and $x_2 = \langle e_1 \otimes e_2 + ve_2 \otimes \beta \rangle$, $u \neq 0 \neq v$. Since the α, e_2, β are linearly independent, we may assume $u\alpha = e_1$ and $v\beta = e_3$. This proves the claim. We call this orbit $\mathbf{o}_{11}$.

$\mathbf{o}_{11}^T$. Replacing $S_{2,3}(\mathbb{F})$ with $S_{3,2}(\mathbb{F})$ in the above, and repeating the argument, we get another orbit $\mathbf{o}_{11}^T$, which is G_1 -equivalent to $\mathbf{o}_{11}$. These can not be H_1 -equivalent, for if they were then $\text{PG}(A_1)$ would be contained in some $\langle S_{2,2}(\mathbb{F}) \rangle$, contradicting our assumption.

The next step in the proof is to consider the remaining orbits of lines which have at least two points of rank two, and no points of rank one. The assumption that such a line $\langle x_1, x_2 \rangle$, where x_1 and x_2 are points of rank two, is not in one of the previous orbits implies that it is not contained in any subspace spanned by a subvariety $S_{2,3}(\mathbb{F})$ or $S_{3,2}(\mathbb{F})$ of $S_{3,3}(\mathbb{F})$, and hence $Q(x_1)$ and

$Q(x_2)$ must intersect in a point, say z . The orbits of such lines depend on the position of the points x_1, x_2, z . We have the following three possibilities: (i) none of x_1, x_2 lies on a secant through z ; (ii) only one of x_1, x_2 lies on a secant through z ; (iii) both x_1 and x_2 lie on a secant through z .

(i) **$\mathbf{o}_{12}$** . In this case we assume w.l.o.g. that $z = \langle e_1 \otimes e_2 \rangle$. Denote the two planes of the Segre variety $S_{3,3}(\mathbb{F})$ that contain z by $\pi(z) = \text{PG}(\mathbb{F}^3 \otimes e_2)$ and $\pi'(z) = \text{PG}(e_1 \otimes \mathbb{F}^3)$. Then the hypotheses imply that for each $i \in \{1, 2\}$, x_i lies on the tangent plane of $Q(x_i)$ at z , and hence x_i is on a secant line $\langle y_i, y'_i \rangle$ with $\langle z, y_i \rangle = \pi(z) \cap Q(x_i)$ and $\langle z, y'_i \rangle = \pi'(z) \cap Q(x_i)$. Summarizing, w.l.o.g. we may assume that $x_i = \langle \alpha_i \otimes e_2 + e_1 \otimes \alpha'_i \rangle$, where both e_1, α_1, α_2 and $e_2, \alpha'_1, \alpha'_2$ are linearly independent. It follows that the line $\langle x_1, x_2 \rangle$ is equivalent to the line $\langle e_1 \otimes e_1 + e_2 \otimes e_2, e_1 \otimes e_3 + e_3 \otimes e_2 \rangle$, under the action induced by (g_1, g_2) , where $g_1 : e_1 \mapsto e_1, \alpha_1 \mapsto e_2, \alpha_2 \mapsto e_3, g_2 : e_2 \mapsto e_2, \alpha'_1 \mapsto e_1, \alpha'_2 \mapsto e_3$. We call this orbit **$\mathbf{o}_{12}$** , and observe that a line in this orbit is a constant rank two line not contained in the span of any subvariety $S_{2,3}(\mathbb{F})$ or $S_{3,2}(\mathbb{F})$.

(ii) **$\mathbf{o}_{13}$** . In this case we assume w.l.o.g. that $z = \langle e_1 \otimes e_2 \rangle$, and that x_2 lies on a secant through z . Then by hypothesis x_1 does not lie on a secant through z , hence, using the same notation as in case (i) x_1 lies on the tangent plane of $Q(x_1)$ at z , and therefore on a secant line $\langle y_1, y'_1 \rangle$ with $\langle z, y_1 \rangle = \pi(z) \cap Q(x_1)$ and $\langle z, y'_1 \rangle = \pi'(z) \cap Q(x_1)$. Continuing the argument as before, we may conclude that w.l.o.g. $x_1 = \langle e_1 \otimes e_1 + e_2 \otimes e_2 \rangle$ and $x_2 = \langle e_1 \otimes e_2 + e_3 \otimes e_3 \rangle$. We denote this orbit by **$\mathbf{o}_{13}$** ; its lines have rank distribution $[0, 2, q-1]$ over $\mathbb{F}_q$.

(iii) **$\mathbf{o}_{14}$** . In this case w.l.o.g. we assume $z = e_2 \otimes e_2$. Then there are $\alpha, \beta, \gamma, \delta$ such that $x_1 = \alpha \otimes \beta + e_2 \otimes e_2$ and $x_2 = e_2 \otimes e_2 + \gamma \otimes \delta$. Since $\langle x_1, x_2 \rangle$ is not contained in any subspace spanned by a subvariety $S_{2,3}(\mathbb{F})$ or $S_{3,2}(\mathbb{F})$ of $S_{3,3}(\mathbb{F})$, we have that both α, e_2, γ and β, e_2, δ are linearly independent. It follows that each such line is equivalent to $\langle x_1, x_2 \rangle$ with $x_1 = e_1 \otimes e_1 + e_2 \otimes e_2$ and $x_2 = e_2 \otimes e_2 + e_3 \otimes e_3$. We call this orbit **$\mathbf{o}_{14}$** ; its lines have rank distribution $[0, 3, q-2]$ over $\mathbb{F}_q$.

We have now dealt with all the orbits on lines which have at least one point of rank one and with all the orbits on lines with at least two points of rank two. So from now on we only need to consider the lines $\text{PG}(A_1)$ with rank distribution $[a, b, c]$ where $a = 0$ and $b \leq 1$.

Case (4): $\text{PG}(A_1)$ is a line, with $a = 0$, $b = 1$; $\mathbf{o}_{15}, \mathbf{o}_{16}$.

If $b = 1$, then let x_1 be a point of rank three and x_2 a point of rank two on $\text{PG}(A_1)$. We distinguish the following two cases: a) there exist two points y_i , $i = 1, 2$ of rank i such that x_1 is on the line $\langle y_1, y_2 \rangle$ and $Q(y_2) = Q(x_2)$; and b) such points do not exist.

a) $\mathbf{o}_{15}$. In this case w.l.o.g. we may assume that $y_1 = \langle e_3 \otimes e_3 \rangle$ and $Q(y_2) = Q(x_2) = \sigma_{3,3}(\langle e_1, e_2 \rangle \times \langle e_1, e_2 \rangle)$. Now projecting the line $\langle x_1, x_2 \rangle$ from y_1 onto $\langle Q(x_2) \rangle$ gives the constant rank two line $\langle x_2, y_2 \rangle$ in $\langle Q(x_2) \rangle$ (cf. orbit o_{10}). Using the transitivity of the stabilizer of $Q(x_2)$ inside the group induced by H_1 on these lines (cf. orbit o_{10}), we obtain one orbit of such lines. We denote this orbit by $\mathbf{o}_{15}$. A representative for the orbit is $\langle e_1 \otimes e_1 + e_2 \otimes e_2 + ue_1 \otimes e_2 + e_3 \otimes e_3, e_1 \otimes e_2 + ve_2 \otimes e_1 \rangle$, with $u, v \in \mathbb{F}$ such that $v\lambda^2 + uv\lambda - 1 \neq 0$ for all $\lambda \in \mathbb{F}$.

If $\mathbb{F} = \mathbb{F}_q$, the orbit o_{15} has rank distribution $[0, 1, q]$. If $\mathbb{F}$ is algebraically closed, this orbit is empty.

b) $\mathbf{o}_{16}$. In this case let $Q(x_2) = \sigma_{3,3}(\ell \times m)$, and suppose $x_1 = \langle e_1 \otimes e_1 + e_2 \otimes e_2 + e_3 \otimes e_3 \rangle$. Then

(**) x_1 can be written as $x_1 = \langle a_1 \otimes b_1 + a_2 \otimes b_2 + a_3 \otimes b_3 \rangle$, if and only if there exists a nonzero constant $k \in \mathbb{F}$ such that $a_i \cdot b_j = 0$ for $i \neq j$ and $a_i \cdot b_i = k$, where $\cdot$ denotes the usual dot product of vectors.

Recall that an element of the stabiliser N of x_1 has the form (g, g^{-1}) , and corresponds to conjugation in $\text{GL}(3, \mathbb{F})$. By Lemma 2.3, the set of elements (g, g^{-1}) in N such that g stabilises a point (line) x in the first factor corresponds to the set of elements (g, g^{-1}) such that g^{-1} stabilises a line (point) in the second factor, and vice versa.

Now assume $\ell = \langle a_1, a_2 \rangle$, and consider the duality from $\text{PG}(\mathbb{F}^3)$ (the first factor in the pre-image of $\sigma_{3,3}$) to $\text{PG}(\mathbb{F}^3)$ (the second factor in the pre-image of $\sigma_{3,3}$) induced by the standard inner product. The dual space of a subspace will be denoted by $\perp$. If m does not pass through $\langle b_3 \rangle := \ell^\perp$, then put $\langle b_1 \rangle = \langle a_2 \rangle^\perp \cap m$, $\langle b_2 \rangle = \langle a_1 \rangle^\perp \cap m$, $\langle a_3 \rangle = m^\perp$. After appropriate scaling we have the necessary conditions (**) to write $x_1 = \langle a_1 \otimes b_1 + a_2 \otimes b_2 + a_3 \otimes b_3 \rangle$ with $Q(x_2) = \sigma_{3,3}(\langle a_1, a_2 \rangle \times \langle b_1, b_2 \rangle)$. This is in contradiction with the hypothesis

(put $y_2 = \langle a_1 \otimes b_1 + a_2 \otimes b_2 \rangle$ and $y_1 = a_3 \otimes b_3$). It follows that m is a line through $\langle b_3 \rangle = \ell^\perp$. Let $m = \langle b_2, b_3 \rangle$ and suppose $x_2 = \langle a_1 \otimes (ab_2 + bb_3) + a_2 \otimes (cb_2 + db_3) \rangle$ for some $a, b, c, d \in \mathbb{F}$. Then $c = 0$ since otherwise $\langle x_1, x_2 \rangle$ would have a rank two point $\langle a_1 \otimes (-cb_1 + ab_2 + bb_3) + (da_2 - ca_3) \otimes b_3 \rangle$ which is different from x_2 , a contradiction. Arguing as before it follows that there is one orbit, which we call o_{16} , satisfying the hypotheses and it is represented by the line $\langle x_1, x_2 \rangle$ with $x_1 = \langle e_1 \otimes e_1 + e_2 \otimes e_2 + e_3 \otimes e_3 \rangle$ and $x_2 = \langle e_1 \otimes e_2 + e_2 \otimes e_3 \rangle$. If $\mathbb{F} = \mathbb{F}_q$, this orbit has rank distribution $[0, 1, q]$, and is denoted by $\mathbf{o}_{16}$.

This completes the classification of lines containing at least one point of rank less than 3. $\square$

Remark 3.2. *The existence of lines in orbits o_{10} and o_{15} depends on the field $\mathbb{F}$. The orbit o_{10} corresponds to the orbit of a nonsingular point in the Segre variety product of three projective lines, see [9]. These in turn correspond to algebraic field extensions of degree 2. If $\mathbb{F}$ is algebraically closed, then there are no such extensions and hence the orbit o_{10} does not occur. The same holds true for the orbit o_{15} (see proof). For any other field $\mathbb{F}$, if we let $\eta(\mathbb{F})$ denote the number of isotopism classes (see [7, Theorem 4.3]) of algebraic field extensions of degree 2 of $\mathbb{F}$ (which may be infinite), then the number of H -orbits on Ω is $14 + 2\eta(\mathbb{F})$.*

3.3 Lines without singular points

The remaining case, that is the case of lines of constant rank 3, is field-dependent, and it will be convenient to utilise the language of matrices, determinants and characteristic polynomials.

Suppose the line $\langle x_1, x_2 \rangle$, $x_i = \langle v_i \rangle$, has rank distribution $[0, 0, q + 1]$, that is, a constant rank 3 line. We may identify this with a 2-dimensional subspace $\langle v_1, v_2 \rangle$ of $M_3(\mathbb{F})$. Then we have that $f(t) := \det(v_1 - tv_2) \neq 0$ for all $t \in \mathbb{F}$. But f is a polynomial of degree 3 over $\mathbb{F}$, and hence if $\mathbb{F}$ is algebraically closed (or real closed), must have a root in $\mathbb{F}$, a contradiction. So, taking into account the results from Sections 3.1 and 3.2, we have shown the following.

Theorem 3.3. *For any algebraically closed (or real closed) field $\mathbb{F}$, there are precisely 18 H -orbits and 15 G -orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$. There are precisely 20 H -orbits and 17 G -orbits of tensors in $\mathbb{R}^2 \otimes \mathbb{R}^3 \otimes \mathbb{R}^3$.*

We continue with the line $\langle x_1, x_2 \rangle$ without any assumptions on the field $\mathbb{F}$. We may assume that $v_1 = e_1 \otimes e_1 + e_2 \otimes e_2 + e_3 \otimes e_3$, which corresponds to the identity matrix. Recall that the group H_1 acts on matrices as $(g, h) : X \mapsto gXh$. The characteristic polynomial of a tensor v is $p_v(t) := \det(v_1 - tv)$.

Now suppose $\langle v_1, v_2 \rangle$ is equivalent to $\langle v_1, v_3 \rangle$, under an element $(g, h) \in H_1$. Then $g(\alpha v_1 + \beta v_2)h = v_1$, and $gh = \gamma v_1 + \delta v_3$, for some $\alpha, \beta, \gamma, \delta \in \mathbb{F}$.

If $\beta = 0$, then $\delta = 0$, and $h = \alpha g^{-1}$. Hence $g\langle v_1, v_2 \rangle g^{-1} = \langle v_1, v_3 \rangle$, and hence $gv_2g^{-1} = \mu v_1 + \nu v_3$ for some $\mu, \nu \in \mathbb{F}$, i.e. v_2 is similar to $\mu v_1 + \nu v_3$. Clearly it is also true that if v_2 is similar to $\mu v_1 + \nu v_3$, then $\langle v_1, v_2 \rangle$ is equivalent to $\langle v_1, v_3 \rangle$.

If $\beta \neq 0$, then $\delta \neq 0$. Then $h = g^{-1}(\gamma v_1 + \delta v_3)$, and $g(\alpha v_1 + \beta v_2)g^{-1}(\gamma v_1 + \delta v_3) = v_1$. Hence $g(\alpha v_1 + \beta v_2)g^{-1} = (\gamma v_1 + \delta v_3)^{-1}$, and so $(\alpha v_1 + \beta v_2)$ is similar to $(\gamma v_1 + \delta v_3)^{-1}$. It is easy to see that the converse is also true, and hence we have shown the following.

Lemma 3.4. *The line $\langle v_1, v_2 \rangle$ is H_1 -equivalent to $\langle v_1, v_3 \rangle$ if and only if one of the following hold:*

- v_2 is similar to $\mu v_1 + \nu v_3$ for some $\mu, \nu \in \mathbb{F}$;
- $(\alpha v_1 + \beta v_2)$ is similar to $(\gamma v_1 + \delta v_3)^{-1}$ for some $\alpha, \beta, \gamma, \delta \in \mathbb{F}$, $\beta\delta \neq 0$.

Now two matrices are similar if and only if they have the same rational canonical form. As v_2 and v_3 both have irreducible characteristic polynomial, their rational canonical forms are the companion matrices of their respective characteristic polynomials.

We recall the definition of the Mobius transformation of a polynomial: if $\phi = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is an invertible matrix, and f a polynomial of degree n , then

$$f^\phi(t) := (ct + d)^n f\left(\frac{at + b}{ct + d}\right)$$

Note that $f^{a\phi} = f^\phi$ for all $a \in \mathbb{F}$, and so we may take ϕ to be an element of $\text{PGL}(2, q)$.

We recall also the following straightforward lemma.

Lemma 3.5. *Let f be a polynomial of degree n , $C(f)$ its companion matrix, and I the identity matrix. Then*

$$p_{\alpha I + \beta C(f)}(t) = \beta^n f\left(\frac{t - \alpha}{\beta}\right) = f^\phi(t)$$

$$p_{C(f)^{-1}}(t) = -t^n f(0)^{-1} f(-t^{-1}) = f^\rho(t),$$

where $\phi = \begin{pmatrix} 1 & -\alpha \\ 0 & \beta \end{pmatrix}$, $\rho = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

This allows us to prove the following lemma.

Lemma 3.6. *Let v_2 and v_3 be two $n \times n$ matrices with irreducible characteristic polynomials f and g respectively. Then $\langle v_1, v_2 \rangle$ is H_1 -equivalent to $\langle v_1, v_3 \rangle$ if and only if there exists some $\phi \in \text{PGL}(2, q)$ such that $g = f^\phi$.*

Proof: We need to consider the two cases from Lemma 3.4. In the first case, v_2 is similar to $\mu v_1 + \nu v_3$ if and only if $f(t) = p_{\mu v_1 + \nu v_3}(t) = g^\phi(t)$, with $\phi = \begin{pmatrix} 1 & -\mu \\ 0 & \nu \end{pmatrix}$.

For the second case, $(\alpha v_1 + \beta v_2)$ is similar to $(\gamma v_1 + \delta v_3)^{-1}$ if and only if

$$p_{\alpha v_1 + \beta v_2}(t) = p_{(\gamma v_1 + \delta v_3)^{-1}}(t),$$

if and only if

$$f^\psi(t) = p_{\gamma v_1 + \delta v_3}^\rho(t) = g^{\chi\rho}(t),$$

with $\psi = \begin{pmatrix} 1 & -\alpha \\ 0 & \beta \end{pmatrix}$, $\rho = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $\chi = \begin{pmatrix} 1 & -\gamma \\ 0 & \delta \end{pmatrix}$, if and only if

$$f = g^{\chi\rho\psi^{-1}}.$$

Now $\chi\rho\psi^{-1} = \frac{1}{\beta} \begin{pmatrix} \gamma\beta & 1 + \gamma\alpha \\ -\delta\beta & \delta\alpha \end{pmatrix}$, and it is clear that every element of $\text{PGL}(2, q)$ can be represented by one of the forms $\begin{pmatrix} 1 & -\mu \\ 0 & \nu \end{pmatrix}$, $\frac{1}{\beta} \begin{pmatrix} \gamma\beta & 1 + \gamma\alpha \\ -\delta\beta & \delta\alpha \end{pmatrix}$, $\nu, \beta, \delta \neq 0$, proving the claim. $\square$

Theorem 3.7. *There is precisely one orbit of tensors in $\mathbb{F}_q^2 \otimes \mathbb{F}_q^3 \otimes \mathbb{F}_q^3$ whose first contraction space is a line without singular points in $\text{PG}(\mathbb{F}_q^3 \otimes \mathbb{F}_q^3)$.*

Proof: It suffices by Lemma 3.6 to show that every two monic irreducible polynomials of degree 3 are related by a Mobius transformation. Let f be a monic irreducible of degree 3. Then the stabiliser of f under $\text{PGL}(2, q)$ has order 3, and hence the size of the orbit of f under this action is $\frac{|\text{PGL}(2, q)|}{3} = \frac{q^3 - q}{3}$, which is precisely the number of monic irreducible polynomials of degree 3, proving the claim. $\square$

Note that there may be an infinite number of orbits for some fields, such as $\mathbb{Q}$.

Remark 3.8. *We remark that in $\text{PG}(\mathbb{F}_q^n \otimes \mathbb{F}_q^n)$ with $n > 3$, there is more than one orbit of lines of constant rank n over the finite field $\mathbb{F}_q$. For example if n is prime, a similar argument shows that there are at least $\frac{q^{n-1}-1}{q^2-1}$ orbits, arising from lines $\langle v_1, v_2 \rangle$ where v_2 has irreducible characteristic polynomial.*

3.4 Canonical forms of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$

In this section we list the canonical forms for the representatives of the G -orbits in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$. The special cases of Theorems 3.9 and 3.10 for the field $\mathbb{F} = \mathbb{F}_2$ were computed in [1].

The previous subsections are summarized in the following theorem.

Theorem 3.9. *If $\mathbb{F}$ is a finite field, then there are precisely 21 H -orbits and 18 G -orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$. For any algebraically closed field $\mathbb{F}$, there are precisely 18 H -orbits and 15 G -orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^3$. There are precisely 20 H -orbits and 17 G -orbits of tensors in $\mathbb{R}^2 \otimes \mathbb{R}^3 \otimes \mathbb{R}^3$.*

Proof: Follows from Theorems 3.1 and 3.7. $\square$

The canonical forms for the 18 G -orbits in the finite field case can be extracted from the proof of the classification of the orbits. If e_1, e_2, e_3 is a basis for $\mathbb{F}^3$ and $e = e_1 \otimes e_1 + e_2 \otimes e_2 + e_3 \otimes e_3$, then the canonical forms are as follows. The third column contains the rank distribution $r_1(A)$ of the first contraction space of a representative A .

Orbit	Canonical form	$r_1(A)$
o_0	0	$[0, 0, 0]$
o_1	$e_1 \otimes e_1 \otimes e_1$	$[1, 0, 0]$
o_2	$e_1 \otimes (e_1 \otimes e_1 + e_2 \otimes e_2)$	$[0, 1, 0]$
o_3	$e_1 \otimes e$	$[0, 0, 1]$
o_4	$e_1 \otimes e_1 \otimes e_1 + e_2 \otimes e_1 \otimes e_2$	$[q + 1, 0, 0]$
o_5	$e_1 \otimes e_1 \otimes e_1 + e_2 \otimes e_2 \otimes e_2$	$[2, q - 1, 0]$
o_6	$e_1 \otimes e_1 \otimes e_1 + e_2 \otimes (e_1 \otimes e_2 + e_2 \otimes e_1)$	$[1, q, 0]$
o_7	$e_1 \otimes e_1 \otimes e_3 + e_2 \otimes (e_1 \otimes e_1 + e_2 \otimes e_2)$	$[1, q, 0]$
o_8	$e_1 \otimes e_1 \otimes e_1 + e_2 \otimes (e_2 \otimes e_2 + e_3 \otimes e_3)$	$[1, 1, q - 1]$
o_9	$e_1 \otimes e_3 \otimes e_1 + e_2 \otimes e$	$[1, 0, q]$
o_{10}	$e_1 \otimes (e_1 \otimes e_1 + e_2 \otimes e_2 + ue_1 \otimes e_2) + e_2 \otimes (e_1 \otimes e_2 + ve_2 \otimes e_1),$ $v\lambda^2 + uv\lambda - 1 \neq 0 \text{ for all } \lambda \in \mathbb{F}$	$[0, q + 1, 0]$
o_{11}	$e_1 \otimes (e_1 \otimes e_1 + e_2 \otimes e_2) + e_2 \otimes (e_1 \otimes e_2 + e_2 \otimes e_3)$	$[0, q + 1, 0]$
o_{12}	$e_1 \otimes (e_1 \otimes e_1 + e_2 \otimes e_2) + e_2 \otimes (e_1 \otimes e_3 + e_3 \otimes e_2)$	$[0, q + 1, 0]$
o_{13}	$e_1 \otimes (e_1 \otimes e_1 + e_2 \otimes e_2) + e_2 \otimes (e_1 \otimes e_2 + e_3 \otimes e_3)$	$[0, 2, q - 1]$
o_{14}	$e_1 \otimes (e_1 \otimes e_1 + e_2 \otimes e_2) + e_2 \otimes (e_2 \otimes e_2 + e_3 \otimes e_3)$	$[0, 3, q - 2]$
o_{15}	$e_1 \otimes (e + ue_1 \otimes e_2) + e_2 \otimes (e_1 \otimes e_2 + ve_2 \otimes e_1),$ $v\lambda^2 + uv\lambda - 1 \neq 0 \text{ for all } \lambda \in \mathbb{F}$	$[0, 1, q]$
o_{16}	$e_1 \otimes e + e_2 \otimes (e_1 \otimes e_2 + e_2 \otimes e_3)$	$[0, 1, q]$
o_{17}	$e_1 \otimes e + e_2 \otimes (e_1 \otimes e_2 + e_2 \otimes e_3 + e_3 \otimes (\alpha e_1 + \beta e_2 + \gamma e_3)),$ $\lambda^3 + \gamma\lambda^2 - \beta\lambda + \alpha \neq 0 \text{ for all } \lambda \in \mathbb{F}$	$[0, 0, q + 1]$

This classification also allows us to classify tensors in $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^3$. Orbits $o_0, o_1, o_2, o_4, o_4^T, o_5, o_6$, and o_{10} are contained in a space isomorphic to $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^2$. The orbits corresponding to o_2 , o_4 and o_4^T are G -equivalent in $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^2$.

Furthermore, orbits o_7 and o_{11} are contained in a space isomorphic to $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^3$. The orbits corresponding to o_2 and o_4 are G -equivalent in $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^3$, but not G -equivalent to o_4^T . This implies the following.

Theorem 3.10. *If $\mathbb{F}$ is a finite or real field, then there are precisely 10 H -orbits and 9 G -orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^3$. For any algebraically closed field $\mathbb{F}$, there are precisely 9 H -orbits and 8 G -orbits of tensors in $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^3$.*

Remark 3.11. *In [12, Table 4], representatives of each G -orbit in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ are given. For the reader's convenience, we include the correspondence between the orbits $o_0, \dots, o_{17}$ and the orbits as listed in [12]. Note that o_2 and*

o_4 are equivalent under G here, while o_{10}, o_{15} and o_{17} are empty.

	o_0	o_1	o_2	o_3	o_4	o_5	o_6	o_7	o_8	o_9	o_{11}	o_{12}	o_{13}	o_{14}	o_{16}
[12]	25	24	23	22	23	20	21	19	15	16	18	17	12	9	13
[15]															

References

- [1] M.R. Bremner; J. Hu. Canonical forms of small tensors over $\mathbb{F}_2$, arxiv.org/abs/1206.5179v1.
- [2] M.R. Bremner; S.G. Stavrou. Canonical forms of $2 \times 2 \times 2$ and $2 \times 2 \times 2 \times 2$ arrays over $\mathbb{F}_2$ and $\mathbb{F}_3$, *Linear and Multilinear Algebra*, 61 (2013), 986-997.
- [3] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.6.2; 2013. (<http://www.gap-system.org>).
- [4] J. Bamberg; A. Betten; P. Cara; J. De Beule; M. Lavrauw; M. Neunhoffer. FinInG - a GAP package for Finite Incidence Geometry, 1.01 2014, (<http://cage.ugent.be/geometry/fining>).
- [5] F. R. Gantmacher, The theory of matrices. Vols. 1, 2, Translated by K. A. Hirsch, Chelsea Publishing Co., New York, 1959.
- [6] J.M. Landsberg. Tensors: Geometry and Applications. 2012. Graduate Studies in Mathematics, 128. American Mathematical Society, Providence, RI, 2012. xx+439 pp. ISBN: 978-0-8218-6907-9.
- [7] M. Lavrauw. Finite semifields and nonsingular tensors. *Des. Codes Cryptogr.* 68 (2013), 205-227.
- [8] M. Lavrauw, O. Polverino; *Finite semifields*. Chapter in *Current research topics in Galois Geometry* (Editors J. De Beule and L. Storme), NOVA Academic Publishers, New York, 2011.
- [9] M. Lavrauw; J. Sheekey. Orbits of the stabiliser group of the Segre variety product of three projective lines. *Finite Fields Appl.* 26 (2014) 1–6.

- [10] M. Lavrauw; J. Sheekey. Aspects of tensor products over finite fields and Galois geometries. Proceedings of the Academy Contact Forum *Galois Geometries and Applications* at the *Royal Flemish Academy of Belgium for Science and the Arts* (October 5, 2012), (2014), S. Nikova, B. Preneel and L. Storme, Eds., pp. 95–102.
- [11] M. Lavrauw; J. Sheekey. Classification of subspaces in $\mathbb{F}^2 \otimes \mathbb{F}^3$ and orbits in $\mathbb{F}^2 \otimes \mathbb{F}^3 \otimes \mathbb{F}^r$. In preparation.
- [12] A. G. Nurmiev, Orbits and invariants of third-order matrices, *Mat. Sb.* 191 (2000) 101-108.
- [13] R. Shaw; N. Gordon; H. Havlicek. Aspects of the Segre variety $S_{1,1,1}(2)$. *Des. Codes Cryptogr.* 62 (2): 225-239, 2012.
- [14] R. M. Thrall, J. H. Chanler. Ternary trilinear forms in the field of complex numbers, *Duke Math. J.*, 4 (1938) 678-690.
- [15] R. M. Thrall. Metabelian groups and trilinear forms, *Amer. J. Math.*, 60 (1938) 383-415.